

学校编码: 10384

分类号\_\_\_\_\_密级\_\_\_\_\_

学号: 200325014

UDC \_\_\_\_\_

厦 门 大 学  
硕 士 学 位 论 文

荧光比率法检测过氧亚硝酸根阴离子  
的研究以及 CdTe 量子点-米托蒽醌复合物  
的形成及其应用研究

Study on the Ratiometric Fluorescence for the Detection of  
Peroxynitrite and Formation of CdTe Quantum Dots - Mitoxantrone  
Conjugate and Its Application

杜 俊 鸥

指导教师姓名: 郭祥群 教授

专 业 名 称: 分 析 化 学

论文提交日期: 2007 年 1 月

论文答辩时间: 2007 年 月

学位授予日期: 2007 年 月

答辩委员会主席: \_\_\_\_\_

评 阅 人: \_\_\_\_\_

2007 年 1 月

**Study on the Ratiometric Fluorescence for the Detection of  
Peroxynitrite and Formation of CdTe Quantum Dots -  
Mitoxantrone Conjugate and Its Application**

A Dissertation Presented

By

DU JUN-OU

Supervisor: Prof. GUO XIANGQUN

Submitted to the Graduate School of Xiamen University

for the Degree of

MASTER OF PHILOSOPHY

Department of Chemistry, Xiamen University

January, 2007

# 厦门大学学位论文原创性声明

兹呈交的学位论文，是本人在导师指导下独立完成的研究成果。本人在论文写作中参考的其他个人或集体的研究成果，均在文中以明确方式标明。本人依法享有和承担由此论文产生的权利和责任。

声明人（签名）：

年 月 日

# 厦门大学学位论文著作权使用声明

本人完全了解厦门大学有关保留、使用学位论文的规定。厦门大学有权保留并向国家主管部门或其指定机构送交论文的纸质版和电子版，有权将学位论文用于非赢利目的的少量复制并允许论文进入学校图书馆被查阅，有权将学位论文的内容编入有关数据库进行检索，有权将学位论文的标题和摘要汇编出版。保密的学位论文在解密后适用本规定。

本学位论文属于

1、保密（ ），在          年解密后适用本授权书。

2、不保密（ ）

（请在以上相应括号内打“√”）

作者签名：                      日期：          年    月    日

导师签名：                      日期：          年    月    日

# 目录

摘要 .....	I
Abstract .....	III

## 第一部分 荧光比率法检测过氧亚硝酸根阴离子的研究

第一章 前言 .....	2
1.1 生物活性自由基 .....	2
1.2 过氧亚硝酸根阴离子的研究进展 .....	2
1.2.1 ONOO <sup>-</sup> 的产生与分解 .....	3
1.2.2 ONOO <sup>-</sup> 的化学及生物化学效应 .....	4
1.2.3 ONOO <sup>-</sup> 与二氧化碳的反应 .....	5
1.2.4 ONOO <sup>-</sup> 的表征方法 .....	6
1.2.4.1 荧光分析法 .....	6
1.2.4.2 化学发光法 .....	8
1.2.4.3 电子自旋共振法 .....	9
1.2.4.4 分光光度法 .....	10
1.2.4.5 自旋标记荧光探针法 .....	11
1.2.4.6 电化学分析法 .....	12
1.3 荧光比率法的原理及应用研究进展 .....	13
1.3.1 基于荧光共振能量转移机理的荧光比率法 .....	13
1.3.2 基于分子内电荷转移机理的荧光比率法 .....	15
1.3.3 基于激发态分子内质子转移机理的荧光比率法 .....	16
1.3.4 基于参比物质的荧光比率法 .....	16
1.3.5 基于激基缔合物形成机理的荧光比率法 .....	17
1.4 论文的提出及研究内容 .....	17
第二章 比率法检测过氧亚硝酸根阴离子的研究 .....	18

第一节 基于铽络合物发光的荧光比率法检测过氧亚硝酸根阴离子的探索 .....	18
2.1.1 引言 .....	18
2.1.2 实验部分 .....	19
2.1.2.1 仪器与试剂 .....	19
2.1.2.2 实验方法 .....	19
2.1.3 结果与讨论 .....	20
2.1.3.1 激发波长和测定波长的选择 .....	20
2.1.3.2 缓冲溶液酸度对体系的影响 .....	21
2.1.3.3 TPA 浓度对 Tb(III)发光强度的影响 .....	21
2.1.3.4 不同缓冲体系对 TPA 敏化 Tb(III)发光和 TPA 羟基化的影响 .....	22
2.1.3.5 TPA—Tb(III)络合物对 ONOO <sup>-</sup> 的响应 .....	22
2.1.4 小结 .....	24
第二节 基于芘激基缔合物形成的荧光比率法检测过氧亚硝酸根阴离子的探索 .....	25
2.2.1 引言 .....	25
2.2.2 实验方法 .....	26
2.2.3 结果与讨论 .....	26
第三节 基于同步荧光比率法检测过氧亚硝酸根阴离子的研究 .....	27
2.3.1 引言 .....	27
2.3.2 实验部分 .....	28
2.3.2.1 仪器和试剂 .....	28
2.3.2.2 实验方法 .....	29
2.3.3 结果与讨论 .....	29
2.3.3.1 反应体系的光谱特性 .....	29
2.3.3.2 $\Delta\lambda$ 的选择 .....	32
2.3.3.3 反应时间对体系荧光强度的影响 .....	32
2.3.3.4 溶液酸度对反应体系荧光强度的影响 .....	33
2.3.3.5 总碳酸盐浓度对体系荧光强度的影响 .....	34
2.3.3.6 L-酪氨酸浓度对体系荧光强度的影响 .....	36
2.3.3.7 狭缝对荧光强度比值的影响 .....	37

2.3.3.8 线性范围, 检测限和精密度 .....	39
2.3.3.9 与常规荧光的比较 .....	41
2.3.3.10 干扰的考察 .....	42
2.3.4 小结 .....	43
<b>第三章 抗癌药米托蒽醌对二氧化氮自由基清除作用的荧光法评价研究 .....</b>	<b>44</b>
3.1 引言 .....	44
3.2 实验部分 .....	45
3.2.1 仪器和试剂 .....	45
3.2.2 实验方法 .....	45
3.3 结果与讨论 .....	46
3.3.1 检测原理与体系的光谱特征 .....	46
3.3.2 MXT 对 $\cdot\text{NO}_2$ 自由基的清除作用 .....	46
3.3.3 机理讨论 .....	48
3.4 小结 .....	48
<b>第四章 结论与展望 .....</b>	<b>49</b>
<b>参考文献 .....</b>	<b>51</b>

## **第二部分 CdTe 量子点-米托蒽醌复合物的形成及其应用研究**

<b>第一章 CdTe 量子点与米托蒽醌复合物的形成及其应用研究 .....</b>	<b>66</b>
1.1 引言 .....	66
1.2 实验部分 .....	67
1.2.1 仪器和试剂 .....	67
1.2.2 实验方法 .....	68
1.2.2.1 量子点的合成 .....	68
1.2.2.2 QD-MXT 复合物的合成 .....	68

1.2.2.3 荧光检测 .....	69
1.3 结果与讨论 .....	69
1.3.1 量子点粒径的表征 .....	69
1.3.2 CdTe 量子点与 MXT 作用的光谱特性 .....	70
1.3.3 MXT 对量子点荧光猝灭的研究 .....	71
1.3.4 QD-MXT 复合物与 DNA 作用的研究 .....	74
1.3.4.1 QD-MXT 复合物与 DNA 作用的光谱特性 .....	74
1.3.4.2 离子强度的影响 .....	76
1.3.4.3 不同核苷和核苷酸与 QD-MXT 复合物相互作用的研究 .....	78
1.3.4.4 QD-MXT 复合物与热变性的 DNA 相互作用的研究 .....	78
1.3.5 反应条件的优化 .....	79
1.3.5.1 QD-MXT 复合物荧光对 DNA 响应的动力学行为 .....	79
1.3.5.2 QD-MXT 复合物的浓度对体系荧光强度的影响 .....	80
1.3.6 标准曲线 .....	81
1.3.7 干扰的考察 .....	82
1.3.8 对合成样品的分析 .....	83
1.4 小结 .....	83
 第二章 结论与展望 .....	 84
参考文献 .....	86



## Content

<b>Abstract in Chinese</b> .....	I
----------------------------------	---

<b>Abstract in English</b> .....	II
----------------------------------	----

<b>Part One: Study on the Ratiometric Fluorescence for the Detection of Peroxynitrite</b> .....	1
---	---

<b>Chapter One: Introduction</b> .....	2
--	---

1.1 Brief Introduction to the Biological Free Radicals.....	2
---	---

1.2 Introduction to the Study of Peroxynitrite.....	2
---	---

1.2.1 Formation and decomposition of peroxynitrite .....	3
--	---

1.2.2 The chemical and biochemical characters of peroxynitrite.....	4
---	---

1.2.3 The reaction between peroxynitrite and carbon dioxide.....	5
--	---

1.2.4 The determination methods for peroxynitrite .....	6
---	---

1.2.4.1 Fluorescent spectrophotometry .....	6
---	---

1.2.4.2 Chemiluminescence .....	8
---------------------------------	---

1.2.4.3 Electronic spin resonance.....	9
--	---

1.2.4.4 Spectrophotometric method .....	10
---	----

1.2.4.5 Spin-labelled fluorophores .....	11
--	----

1.2.4.6 Electrochemistry .....	12
--------------------------------	----

1.3 Introduction to the Principles and Applications of Ratiometric Fluorescence.....	13
--	----

1.3.1 The ratiometric fluorescence based on FRET.....	13
---	----

1.3.2 The ratiometric fluorescence based on ICT.....	15
--	----

1.3.3 The ratiometric fluorescence based on ESIPT.....	16
--	----

1.3.4 The ratiometric fluorescence based on inner reference.....	16
--	----

1.3.5 The ratiometric fluorescence based on the formation of excimer .....	17
--	----

1.4 Design of the Experimentation .....	17
---	----

<b>Chapter Two: Study on the Analytical Methodology for Peroxynitrite</b>	<b>18</b>
<b>Section One: The Ratiometric Fluorescence for Detection of Peroxynitrite Based on Terbium(III) Complex</b>	<b>18</b>
2.1.1 Introduction	18
2.1.2 Experimentation	19
2.1.2.1 Instruments and reagents	19
2.1.2.2 Methods	19
2.1.3 Results and Discussion	20
2.1.3.1 Choice of excitation and emission wavelength	20
2.1.3.2 Effect of pH	21
2.1.3.3 Effect of TPA concentration on the chemiluminescence of Tb(III)	21
2.1.3.4 Comparison of fluorescence response of Tb(III) sensitized by TPA and hydrolyation of TPA in different buffers	22
2.1.3.5 The response of the complex of TPA—Tb(III) to peroxynitrite	22
2.1.4 Brief Summary	24
<b>Section Two: The Ratiometric Fluorescence for Detection of Peroxynitrite Based on the Formation of Excimer of Pyrene</b>	<b>25</b>
2.2.1 Introduction	25
2.2.2 Experimentation	26
2.2.3 Results and Discussion	26
<b>Section Three: The Ratiometric Fluorescence for Detection of Peroxynitrite Based on Synchronous Ratiometric Fluorimetric Spectrum</b>	<b>27</b>
2.3.1 Introduction	27
2.3.2 Experimentation	28
2.3.2.1 Instruments and reagents	28
2.3.2.2 Methods	29
2.3.3 Results and Discussion	29
2.3.3.1 The spectrum of the system	29
2.3.3.2 Choice of $\Delta\lambda$	32
2.3.3.3 Effect of reaction time	32

2.3.3.4 Effect of pH	33
2.3.3.5 Effect of total carbonate concentration	34
2.3.3.6 Effect of L-tyrosine concentration	36
2.3.3.7 Effect of slit	37
2.3.3.8 Calibration curve, sensitivity	39
2.3.3.9 Compared with fluorescence	41
2.3.3.10 Effects of foreign substances	42
2.3.4 Brief Summary	43
<b>Chapter Three: Evaluation of the Scavenging Effect of MXT to <math>\cdot\text{NO}_2</math> by Fluorescence Spectrophotometry</b>	<b>44</b>
3.1 Introduction	44
3.2 Experimentation	45
3.2.1 Instruments and reagents	45
3.2.2 Methods	45
3.3 Results and Discussion	46
3.3.1 Principle of the method and fluorescence spectra	46
3.3.2 Scavenging $\cdot\text{NO}_2$ by MXT	46
3.3.3 Mechanism	48
3.4 Brief Summary	48
<b>Chapter Four: Conclusion and Perspectives</b>	<b>49</b>
<b>Reference</b>	<b>51</b>

## Part Two: Formation of Quantum Dots - Mitoxantrone Conjugate and Its Application

### Chapter One: Study on the Formation of Quantum Dots - Mitoxantrone Conjugate

<b>and Its Application</b> .....	<b>65</b>
1.1 Introduction .....	66
1.2 Experimentation .....	67
1.2.1 Instruments and reagents .....	67
1.2.2 Methods .....	68
1.2.2.1 Synthesis of CdTe QDs .....	68
1.2.2.2 Conjugation of QDs with MXT .....	68
1.2.2.3 Fluorescence determination .....	69
1.3 Results and Discussion .....	69
1.3.1 Particles size determination .....	69
1.3.2 Spectra characterization of the interaction between CdTe QDs and MXT .....	70
1.3.3 Fluorescence quenching of CdTe QDs by MXT .....	71
1.3.4 Study on the interaction between QD-MXT conjugate and DNA .....	74
1.3.4.1 Preliminary optical characterization .....	74
1.3.4.2 Effect of ionic strength .....	76
1.3.4.3 Mutual effect between different nucleoside or nucleic acid and QD-MXT conjugate .....	78
1.3.4.4 Mutual effect between denatured DNA and QD-MXT conjugate .....	78
1.3.5 Optimized assay conditions .....	79
1.3.5.1 Kinetic characteristics .....	79
1.3.5.2 Optimum amounts of QD-MXT conjugate .....	80
1.3.6 Calibration curves and sensitivity .....	81
1.3.7 Tolerance of foreign substances .....	82
1.3.8 Analytical results of DNA in synthetic samples .....	83
1.4 Brief Summary .....	83
<b>Chapter Two: Conclusion and Perspectives</b> .....	<b>84</b>
<b>Reference</b> .....	<b>86</b>

## 摘 要

本论文研究内容分为两部分。第一部分为荧光比率法表征过氧亚硝酸根阴离子 ( $\text{ONOO}^-$ ) 的方法研究, 及基于此建立抗癌药物米托蒽醌抑制二氧化氮自由基作用评价的比率荧光法研究; 第二部分为 CdTe 量子点-米托蒽醌复合物的形成及应用于核酸检测的初步研究。第二部分内容是基于课题组现行研究课题的新的发展方向, 为实验中观察到的现象的拓展。

第一部分第一章首先简要介绍了活性氧自由基的定义及研究意义; 综述了  $\text{ONOO}^-$  的形成、分解、性质及其表征方法; 系统阐述了荧光比率法的检测原理和应用的研究进展, 并在此基础上提出了本论文第一部分的研究设想, 引用参考文献 160 篇。

第一部分第二章对荧光比率法检测过氧亚硝酸根阴离子的方法进行了研究, 该章共分为三节。

第一节: 基于铽络合物发光的荧光比率法检测  $\text{ONOO}^-$  的研究。 $\text{ONOO}^-$  在弱酸性介质中易质子化生成其共轭酸 ( $\text{ONOOH}$ ),  $\text{ONOOH}$  均裂生成  $\cdot\text{OH}$ 。 $\cdot\text{OH}$  作用于对苯二甲酸发生芳环羟基化反应, 生成强荧光物质邻羟基对苯二甲酸。对苯二甲酸具有敏化铽发光的特性, 羟基化导致其敏化活性消失。因此, 在对苯二甲酸-铽敏化发光体系中加入  $\text{ONOO}^-$ , 将观察到铽的特征发射受到抑制, 同时伴随着邻羟基对苯二甲酸的特征发射峰的出现, 且随着  $\text{ONOO}^-$  浓度的加大, 430 nm 处的邻羟基对苯二甲酸特征发射与 545 nm 处的铽特征发射之比值呈增大的趋势, 从而可以利用这两个峰的强度变化比来表征  $\text{ONOO}^-$ 。

第二节: 基于蒽激基缔合物形成的荧光比率法检测  $\text{ONOO}^-$  的初步研究。该项研究的基本思路是发展基于蒽的二聚体-单体平衡的、以  $\text{ONOO}^-$  为目标分子的比率型荧光探针, 即以蒽分子标记酪胺, 当酪胺与目标分子  $\text{ONOO}^-$  作用时易发生二聚, 酪胺的二聚, 拉近了分别标记于两个酪胺分子上的蒽分子, 两个蒽分子由于距离接近而出现激基缔合物发射, 与之同时, 伴随着单体发射的荧光峰减弱, 激基缔合物态与单体态发射的荧光强度之比  $I_E/I_M$  应与目标分子  $\text{ONOO}^-$  的浓度相关, 以此为基础发展建立基于荧光比率原理的  $\text{ONOO}^-$  荧光分析法。这部分实验只进行了初步的探索, 尚未达到预期的效果。

第三节: 基于酪氨酸二聚的同步荧光法比率法检测  $\text{ONOO}^-$  的研究。酪氨酸与目标

分子  $\text{ONOO}^-$  作用时, 如果控制酪氨酸的浓度远大于  $\text{ONOO}^-$ , 可得到单体发射恒定不变而二聚体的发射则随目标分子的浓度的增大而增强的同步发射光谱特征。采用同步荧光扫描, 以酪氨酸自身的荧光作内参比, 二聚酪氨酸的荧光强度与酪氨酸自身的荧光强度的比值与  $\text{ONOO}^-$  的浓度在一定范围内呈线性相关, 且能不受仪器测试参数改变影响, 从而建立了一种基于内参比的同步荧光比率检测  $\text{ONOO}^-$  的方法。

第一部分第三章是在第二章第三节的实验结果的基础上, 建立了用于评价抗癌药米托蒽醌清除二氧化氮自由基效力的荧光法。二氧化碳催化  $\text{ONOO}^-$  产生  $\cdot\text{NO}_2$ ,  $\cdot\text{NO}_2$  诱导强荧光酪氨酸二聚体形成。米托蒽醌具有抗氧化活性, 当体系中引入米托蒽醌时, 米托蒽醌与酪氨酸竞争  $\cdot\text{NO}_2$ , 抑制酪氨酸二聚, 表现为二聚体荧光强度降低。据此建立了一种间接表征米托蒽醌抗氧化效力的方法。

第一部分第四章是对第一部分的研究工作的总结, 指出工作的不足, 并对研究工作的前景提出了设想。

第二部分第一章首先简要介绍了量子点及其复合物研究意义, 并在此基础上提出了研究的设想。然后根据文献合成了水溶性的巯基乙酸包裹的 CdTe 量子点, 进而依据静电作用原理, 制备了量子点-米托蒽醌 (QD-MXT) 复合物。与 CdTe 量子点具有强发光特性形成强烈反差的是该复合物中量子点的荧光几乎完全猝灭, 为一非荧光复合物。当 QD-MXT 复合物作用于 DNA 后, 由于 DNA 与米托蒽醌发生分子间作用, 导致量子点荧光回复。基于对光谱性质的分析, 就复合物发光猝灭机制与 DNA 响应机制进行了探讨。对应用 QD-MXT 复合物表征 DNA 进行了初步研究。

第二部分第二章是对第二部分的研究工作的总结, 指出工作的不足, 并对研究工作的前景提出了设想。

关键词: 过氧亚硝酸根阴离子; 荧光比率法; 酪氨酸; 量子点; 复合物; DNA

## Abstract

The dissertation consists of two parts. The first part is focused on the study on the analytical methodology based on ratiometric fluorescence for the detection of peroxynitrite ( $\text{ONOO}^-$ ) and evaluation of the scavenging effect of mitoxantrone to  $\cdot\text{NO}_2$ . The second one presents some preliminary experimental results related to CdTe Quantum Dots - mitoxantrone conjugate and its application in determination of nucleic acids.

In chapter 1 of part one, a brief review on the biological active species,  $\text{ONOO}^-$ , is given, concerning its formation, decomposition, reactivity, and its biological significance, as well as the methods for the determination of peroxynitrite that have been developed and used in analytical science and bioscience. A systematic review on the theory and application of ratiometric fluorescence are also presented. 160 references are cited. The research plan for part one of the present dissertation is given.

In chapter 2 of part one, the results of a series of studies on the analytical methodology based on ratiometric fluorescence for the detection of peroxynitrite are presented. This chapter contains three sections.

In section one, a study on ratiometric determination of  $\text{ONOO}^-$  using Terbium(III) complex as a fluorescent probe is presented.  $\text{ONOO}^-$  decomposes to free radical of nitrite ( $\cdot\text{NO}_2$ ) and hydroxyl radical ( $\cdot\text{OH}$ ) on protonation.  $\cdot\text{OH}$  would react rapidly with terephthalic acid and transfer the latter to a high fluorescence compound, 2-hydroxyl-terephthalic acid. Terephthalic acid would sensitize the luminescence of Tb(III) through an intramolecular energy transfer manner on coordinating with Tb(III). The formation of 2-hydroxyl-terephthalic acid could greatly attenuate the sensitization of Tb(III)'s luminescence, thus, an increase in the fluorescence intensity of 2-hydroxyl-terephthalic acid at 430 nm and a decrease of the fluorescence intensity of Tb(III) at 545 nm would be observed in the presence of  $\text{ONOO}^-$ . The intensity ratio,  $F_{430}/F_{545}$ , was proportional to the concentration of the hydroxyl radical, and thus to the concentration of peroxynitrite.

In section two, a ratiometric fluorimetric method for detection of peroxynitrite based

on the formation of excimer of pyrene was explored. Tyramine was labeled with pyrene. When dityramine was formed due to reaction tyramine with peroxynitrite, excimer of pyrene would form as a result of the short distance between the two pyrene molecules. The increase in fluorescence of excimer would be observed and, at the same time, the decrease in fluorescence of monomer of pyrene would be measured. The ratio of the fluorescence intensity of monomer ( $I_M$ ) versus the fluorescence intensity of excimer ( $I_E$ ) would have a good response to peroxynitrite. This hypothesis, however, the expected results have not been gotten yet.

In section three, a ratiometric synchronous fluorimetric method for detection of peroxynitrite was developed based on the formation of dityrosine. Tyrosine acted as inner reference. The ratio of the fluorescence intensity of dityrosine versus the fluorescence intensity of tyrosine was not affected by instrument parameter and had a good response to peroxynitrite by synchronous fluorescence spectrometry.

In chapter 3 of part one, based on the result obtained in section three of chapter 2, a new method was established for evaluation of the scavenging effect of mitoxantrone to  $\cdot\text{NO}_2$  by fluorescence spectrophotometry. The transformation of tyrosine to dityrosine is mediated by  $\cdot\text{NO}_2$  produced from  $\text{ONOO}^-$  and catalyzed by  $\text{CO}_2$ . The production of dityrosine was attenuated in the presence of mitoxantrone due to the scavenging action of mitoxantrone on  $\cdot\text{NO}_2$  and, thus the fluorescence produced by dityrosine was reduced. The change of fluorescence had a good response to mitoxantrone.

In chapter 4 of part one, the research work of part one was summarized, and the expectation and limitation of this research were given.

In chapter 1 of part two, the brief introduction concerning quantum dots and its conjugate are given. Water-soluble CdTe quantum dots were prepared and conjugated with mitoxantrone through electrostatic interaction. Upon this reaction, the fluorescence of QD was quenched. When DNA was added, the fluorescence of QD was increased. A possible mechanism of the interaction was proposed. Thus, a novel approach was developed for detection of DNA.

In chapter 2 of part two, the research work of part one was summarized, and the



Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to [etd@xmu.edu.cn](mailto:etd@xmu.edu.cn) for delivery details.

厦门大学博硕士论文摘要库